

^{10}B in the form of Boric Acid for nuclear industry



- B-10 enriched Boric Acid for use as a chemical shim for excess neutron absorption in the primary circuit of PWRs using high burnup or MOX fuel cores.
- In these PWRs neutron absorption of natural Boric Acid is insufficient. Solubility of Boric Acid limits the boron concentration in primary circuit water to be used as neutron poison.
- Today more and more enriched Boric Acid (>95% enriched in ^{10}B) is used by NPPs.

NUKEM Isotopes GmbH

Rodenbacher Str. 47, 63755 Alzenau, Germany, T: +49 (0)6023 9474-800, F: +49 (0)6023 9474-813
E: info@nukemisotopes.de, I: www.nukem-isotopes.com

¹⁰B in the form of Boric Acid

Specification

Physical properties:

- Material ¹⁰B – Boron-10 in the form of crystalline Boric Acid
- Enrichment ¹⁰B ≥ 96at%

Chemical Properties:

- Form B(OH)₃
- Purity ≥ 99,95 wt%

Impurities in ppm

As	≤ 0,5
Ca	≤ 1
Cl	≤ 5
F	≤ 1
Na	≤ 5
Pb	≤ 1
PO ₄	≤ 1
SO ₄	≤ 2
Insolubles in H ₂ O	≤ 20

NUKEM Isotopes GmbH

Rodenbacher Str. 47, 63755 Alzenau, Germany, T: +49 (0)6023 9474-800, F: +49 (0)6023 9474-813
E: info@nukemisotopes.de, I: www.nukem-isotopes.com